Thromboembolic disease: a geriatric syndrome

Flavia Galdi, Claudio Pedone, Raffaele Antonelli Incalzi

Geriatric Area, Università Campus Bio-Medico, Rome, Italy

The incidence of thrombotic cardiovascular disease increases with age. Several potential reasons can explain the association between thrombosis risk and age, such as the high prevalence of predisposing illnesses among older people and age-related changes in hemostasis pathways, including endothelial dysfunction, alterations in platelet function, coagulation and fibrinolytic factors, that contribute to create a hypercoagulability of the blood. Furthermore, some components of this procoagulant status may also support the inflammatory processes that characterize advancing age, promoting an increased risk of a thrombo-inflammatory disease. However, despite the increased thrombotic risk make the older patient a theoretical target for anticoagulant therapy, the concomitant risk of major bleeding make the use of this drugs problematic, with consequent underuse.

This review summarizes the physiological changes of hemostasis pathways in older people in the context of thromboembolic disease as a geriatric syndrome. The balance between thrombotic and hemorrhagic risk and therefore the therapeutic efficacy of anticoagulant medications are also discussed.

Key words: thrombosis, aging, inflammaging, hypercoagulability, thromboprophylaxis

Received: February 20, 2020

Accepted: April 27, 2020

Correspondence

Flavia Galdi

Geriatric Area, Università Campus Bio-Medico di
Roma, via Alvaro del Portillo 21
00128 Rome, Italy. Tel.: +39 06 225411350.
Fax: +39 06 22541456.
E-mail: f.galdi@unicampus.it

Conflict of interest

The Authors declare no conflict of interest

How to cite this article: Galdi F, Pedone C, Antonelli Incalzi R. Thromboembolic disease: a geriatric syndrome. *Journal of Gerontology and Geriatrics* 2021;69:68-78. <https://doi.org/10.36150/2499-6564-437>

EPIDEMIOLOGY

The incidence of thrombotic cardiovascular disease (including ischemic heart disease, peripheral vascular disease, stroke and venous thromboembolism [VTE]) increases with age ¹. The average age at first myocardial infarction is 65.6 years for males and 72.0 years for females ², and coronary artery disease (CAD) is the principal cause of death in adults aged 75 and older, with more than 80% of all CAD-related deaths occurring in this population ³. In addition, approximately 50% of all strokes occurring in people over age 75 ⁴. The incidence of a first VTE episode, including deep vein thrombosis and pulmonary embolism, increases from < 1 case per 1000 person-years in persons aged < 50 years to 6-8 per 1000 person-years in persons aged > 80 years ⁵ and data on mortality are limited. In particular, pulmonary embolism (PE) represents an increasing proportion of the manifestations of VTE, with an incidence that increases from 120 in 100,000 persons per year in the 65 to 69 years age group to more than 700 in 100,000 subjects per year in the 85 years and older group ⁶. The prognosis of the VTE is very poor in older patients with an overall 30-day mortality that increases from 4% in patients aged < 65 years to 11% in patients aged ≥ 65 years and remains higher after one year (14% vs 31%) ⁷. In the first 3 months following VTE, the most common causes of death in older subjects are cancer (38%),

© Copyright by Società Italiana di Gerontologia e Geriatria (SIGG)

OPEN ACCESS

This is an open access article distributed in accordance with the CC-BY-NC-ND (Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International) license. The article can be used by giving appropriate credit and mentioning the license, but only for non-commercial purposes and only in the original version. For further information: <https://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>

pulmonary embolism (26%), infections (13%) and bleeding complications (6%)⁸.

There are several potential explanations for this association between thrombosis risk and age, first of all the common occurrence of predisposing illnesses among older people. Moreover, the risk factors for arterial and venous thromboembolism are more prevalent in older subjects⁹ and they can be distinguished in modifiable and/or temporary risk factors and non-modifiable risk factors¹⁰ (Tab. I). In addition to the increased level of several plasma components, age-related endothelial dysfunction and alterations in platelet function may contribute to the increased risk of VTE in the elderly⁹.

The aim of this review is to summarize the physiological changes of hemostasis pathways in older people in the context of thromboembolic disease as a geriatric syndrome. In the light of these data, the balance between thrombotic and hemorrhagic risk and therefore the therapeutic efficacy of antithrombotic medications are also discussed.

HEMOSTASIS PATHWAYS AND AGING

There are two main components of hemostasis: primary hemostasis, that refers to platelet aggregation and platelet plug formation, and secondary hemostasis, that refers to the deposition of insoluble fibrin, which is generated by the proteolytic coagulation cascade. These two processes happen simultaneously and are mechanistically intertwined.

Platelets are activated in a multifaceted process, and as a result they adhere to the site of injury and to each other, promoting a developing platelet thrombus. This platelet plug is also stabilized by deposition of insoluble fibrin generated by the coagulation cascade.

When the vascular system is injured tissue factor (TF), a transmembrane glycoprotein, functions as the principal

initiator of the coagulation cascade. TF activates factor VIIa creating an enzymatic complex that activate factor X and factor IX. Factor IXa also activates factor X, in the presence of its cofactor factor VIIIa. Factor Xa, also in the presence of its cofactor factor Va, then activates prothrombin to generate thrombin (IIa) that cleaves soluble fibrinogen to generate insoluble fibrin that forms a crosslinked fibrin mesh at the site of an injury. Fibrin generation occurs simultaneously to platelet aggregation. Thrombin also plays an important role in down regulation of the coagulation cascade by binding to thrombomodulin on endothelial cells and then activating protein C (APC). The activated protein C anticoagulant system, with the cofactor protein S, is important for the down regulation of the coagulation cascade, inactivating the procoagulant cofactors VIIIa and Va. The coagulation cascade is also down-regulated by other physiological inhibitors of blood coagulation: antithrombin, that inhibits thrombin and factor Xa as well as factor IXa and factor XIa; heparin cofactor II (thrombin inhibitor) and tissue factor pathway inhibitor (TFPI), that inhibits factor Xa.

The fibrinolytic system acts to dissolve and remove the clot formed from the injured tissue. The end product of this pathway, plasmin, is a potent enzyme that cleaves fibrin and this generates fibrin degradation products (FDPs), including D-dimer¹¹.

The main age-related changes of the vascular system can be considered within the context of the classic Virchow triad (abnormal vessel wall, abnormal blood flow, and abnormal blood constituents). In addition, there are several changes in the hemostatic system, including coagulation and fibrinolytic factors¹², that contribute to create a hypercoagulability of the blood with an increased incidence of thrombosis in the older people. Furthermore, while changes in hemostasis pathways commonly occur in the general population with aging, some components of this procoagulant status may also support the inflammatory processes that characterize advancing age, since markers such as factor VIII and fibrinogen, are acute-phase proteins¹³ the plasma concentrations of many coagulation factors (e.g., fibrinogen, factor [F] V, FVII, FVIII, and FIX (Fig. 1).

INFLAMMATION, THROMBOSIS AND AGING

Older people are frequently affected by multiple morbidities that are established prothrombotic risk factors, such as congestive heart failure, chronic obstructive disease, diabetes, cancer, chronic venous insufficiency¹⁴. Advanced age is also associated with an increase of inflammatory state that may be another important stimulus for thrombus formation, but the causal pathways and molecular mechanisms that connect inflammaging

Table I. Arterial and venous thromboembolism risk factors.

Modifiable and/or temporary risk factors	Non-modifiable risk factors
Obesity	Age
Elevated homocysteine	Genetic factors such as a deficiency of endogenous anticoagulants (antithrombin, protein C and S)
Hospitalization	Increased concentration of procoagulant proteins (fibrinogen, factors V, VII, VIII, IX, XII, D-dimer, VWF, thrombin)
Cancer	
Surgery	
Trauma	
Immobilization	
Travel

Figure 1 Age-related changes in hemostasis pathway.

and chronic diseases are not well understood. Two major hypotheses concerning the involvement of chronic inflammation in aging have been proposed: molecular inflammation and inflammaging¹⁵. The molecular inflammation hypothesis is based on molecular changes in inflammation-related transcription factors and in the expression levels of their target genes. The hypothesis states that these changes are the mechanism underlying the aging process and age-related diseases^{16,17}. Prolonged oxidative stress and compromised antioxidant defense systems during aging are blamed for increased reactive species (RS), including reactive oxygen species (ROS), reactive nitrogen species (RNS), and reactive lipid aldehydes¹⁸. Although young organisms have a well-functioning antioxidant system to maintain redox balance, the age-related decline in the anti-oxidant defense leads to failure to maintain redox homeostasis, with activation of various proinflammatory signaling pathways. The Inflammaging hypothesis is based on the commonly observed age-associated increase in pro-inflammatory cytokines¹⁹. This concept states that activation of the innate immune system in older individuals leads to a dysregulation in inflammation that impairs the ability to initiate an efficient innate and adaptive immune program in responses to antigens or environmental stimuli²⁰. According to this hypothesis,

these alterations in the immune system contribute to the development of overt organ-specific inflammatory diseases such as atherosclerosis, hypercoagulability, Alzheimer's disease, and diabetes²¹. Clinically, inflammaging is characterized by increased blood levels of several inflammatory biomarkers, including c-reactive protein (CRP), IL-6, IL-18 and tumor necrosis factor- α (TNF- α). In particular, IL-6 is a critical factor in the acute phase of inflammation and also upregulates the synthesis of hemostatic factors, such as fibrinogen²². IL-6 may also directly activate platelets, leading to platelet aggregation with a faster thrombus formation²³. Another element that may promote thrombo-inflammatory responses is an amplified platelet-monocyte interactions due to monocyte phenotype and function that are different in older adults than in younger adults²⁴. In healthy individuals, about 95% of circulating monocytes display the cell surface antigen CD14 and, the remaining smaller fraction of monocytes express CD16, with a pattern CD14^{high}/CD16^{low}. It has been demonstrated that acute systemic inflammation²⁵, but also aging, are associated with a shift toward proinflammatory monocytes pattern CD14^{low}/CD16^{high}²⁴. For example, Seidler et al. showed in a cohort of 181 healthy adults aged 18 to 88, that pattern of CD14^{low}/CD16^{high} monocyte counts increased with age and

demonstrated altered surface protein and chemokine receptor expression²⁶. Also Sadeghi et al. showed in a smaller study of nursing home residents that CD14low/CD16high monocytes produced higher levels of proinflammatory cytokines²⁷.

CHANGES IN VASCULAR SYSTEM AND PLATELETS

The peculiar aspects that characterize the aging of the vascular system include changes in the microcirculation, loss of vein structural integrity, stiffness and dilation of the arteries due to the degeneration of elastic fibers and the increase in collagen content²⁸ and gradual development of endothelial dysfunction. All these changes can increase the risk for thrombosis in older people. The aged blood vessels also express less endothelial nitric oxide (NO) synthase²⁹, resulting in less NO production³⁰. Decreased NO production may contribute to increased platelet activation and arterial thrombosis³¹ as well as enhanced atherogenesis³².

Age-related changes in platelet count and function have also been documented. Comparing platelet count of a large population of older people with that of children, Balduini et al.³³ showed a reduction of 35% in males and 25% in females. A correlation between platelet count and age was also found in a large study including 12,142 American subjects showing an average decrease of 7×10^3 platelets/ μ L and of 18×10^3 platelets/ μ L in individuals in the 60–69 year age group and 69 to 90 years, respectively, compared with those in the 20–59 years age group³⁴. This phenomenon of age-related changes in platelet count has no explanation yet. It has been suggested a reduction of stem cell reserve in older people or a survival advantage of subject with lower platelet count in reaching older age³⁴. In contrast, with advancing age there is a platelet hyperactivity: bleeding time, a measure of platelet aggregation responses, decreases significantly denoting a greater aggregation with a faster thrombus formation³⁵. The mechanisms of this age-related platelet hyperactivity is unclear. Bastyr et al. have tested the hypothesis that the increase of platelet activity with age can be associated with a higher content of platelet phospholipids, suggesting an age-related alteration in platelet transmembrane signaling or second messenger accumulation³⁶. Markers of platelet activation, β -thromboglobulin (a protein stored in platelet granules) and platelet factor 4 (a key platelet receptor), are significantly elevated in older compared with younger people³⁷. Furthermore, with advancing age there is a progressive decline of glutathione peroxidase-3 (GPx3), an antioxidant enzyme, contributing to vascular damage and endothelial dysfunction³⁸. It has been shown that a

deficiency of GPx3, increases platelet activation with a consequent high risk of thrombotic events^{39,40}.

Moreover, with advancing age platelets are less susceptible to inhibition by prostacyclin and have reduced surface prostacyclin receptors density²⁴. Thus, altered activity of platelets contribute to increase the risk of thrombotic disorders in older subjects.

CHANGES IN PROCOAGULANT, ANTICOAGULANT AND FIBRINOLYTIC SYSTEM PROTEINS

Plasma concentrations of several coagulation proteins, including fibrinogen, factors V, VII, VIII, IX and XII, increase with advancing aging¹³. In contrast, concentrations of FX and prothrombin do not change significantly throughout the lifetime⁴¹.

Elevated levels of fibrinogen have been described as a recognized risk factor for thrombotic disorders⁴², even if the mechanism by which fibrinogen contribute to increase cardiovascular risk have not been elucidated. In particular, fibrinogen levels have been shown to increase with age. Kannel et al. demonstrated in the Framingham study that the mean plasma level of fibrinogen increased from about 280 mg/dL in individuals aged 47–54 years to over 300 mg/dL in individuals aged 65–79 years⁴². In a population study, people aged 53–64 years showed significantly higher levels (300 mg/dL) than subjects 20 years old (250 mg/dL)⁴³. Thus, an increment of plasma fibrinogen level by 10 mg/dL for each decade can be expected in healthy subjects¹³.

Factor VIII progressively increase with age, reaching a mean of over 200 U/dL in the seventh decade of life⁴⁴, and it is associated with an approximately fivefold increased risk of venous thrombosis when exceeding 150 U/dL⁴⁵.

Ofori et al. demonstrated age-related changes in FVII proteolysis in normal individuals with a consequent increase in the concentrations of activated FVII, FVII zymogen, total FVII-related proteins, tissue factor pathway inhibitor (TFPI), and prothrombin fragments 1 + 2⁴⁶. Thus, the authors hypothesized that these could lead to increased thrombin generation in vivo, with a possible explanation of the high thrombotic risk in older people. Furthermore, FVII has been described as an important risk factor for cardiovascular events in the Prospective Cardiovascular Münster study⁴⁷, in particular when associated with additional cardiovascular risk factors, such as smoking, myocardial infarction events in family, angina pectoris, high levels of fibrinogen, total cholesterol, LDL cholesterol, and triglycerides, and a low level of HDL cholesterol. In contrast, the Cardiovascular Health Study (CHS), which analyzes 5888 white and African American men and women aged ≥ 65 years,

found an association with cardiovascular events for high levels of fibrinogen and FVIII, but not of FVII. It is likely that methodological differences between the studies might explain this discrepancy ⁴⁸.

Among prothrombotic factors, it has been found that also levels of von Willebrand factor (VWF) increases with aging contributing to higher incidence of thromboembolic disease in older people ⁶. Moreover, it has been showed that ADAMTS13, a metalloproteinase that cleaves VWF preventing its accumulation, does not increase in a parallel fashion, with a consequent additional shift toward the prothrombotic state ⁴⁹.

As far as anticoagulant factors are concerned, the available evidences about their changes with age are contrasting. Favaloro et al. showed small but significant age-related increases in protein C activity, and in total and free protein S, but also a reduction in plasma concentration of antithrombin ⁵⁰. In contrast, Bauer et al found no change in antithrombin and protein C levels with aging in men aged 21 to 81 years ⁵¹.

With respect to the fibrinolytic system, it has been described a reduction in plasminogen levels in subjects aged 75 years or older ⁵², but another study reported that plasminogen levels decreased slightly with age in women but not in men ⁵³. In contrast, plasminogen activator inhibitor (PAI)-1, which is the major inhibitor of fibrinolysis, increased with advancing age ¹⁴. In particular, PAI-1 is an acute phase reactant influenced by several cytokines, including IL-6, TNF- α , TGF- β and hormones, and its expression is related to stress-induced thrombosis in older people. Indeed, Yamamoto et al. demonstrate in aged mice that stress causes a thrombotic tendency in an age-associated manner, and this result correlates with the large induction of PAI-1 expression ⁵⁴.

THE PARADOX OF HYPERCOAGULABILITY IN OLDER POPULATION

While being at first glance a detrimental effect of aging, age-related changes in the coagulation system may also have beneficial and compensatory effects, such as against tumor growth and metastasis ⁶. In fact, several coagulation products act as inhibitors of angiogenesis ^{55,56}. These include domain 5 of high-molecular-weight kininogen, a cleaved form of antithrombin, prothrombin fragment 1+2 (F1+2), and angiostatin, a derivative of plasminogen. Platelet products, including PF4, PAI-1, and thrombospondin-1, are also inhibitors of angiogenesis ⁵⁶. Thus, a higher propensity for thrombosis, although potentially detrimental in patients with atherosclerosis, may confer a survival advantage by inhibiting tumor-associated angiogenesis in the elderly population.

THROMBOPROPHYLAXIS AND RISK OF BLEEDING IN OLDER POPULATION

The increased thrombotic risk make the older patient a theoretical target for anticoagulant therapy, but the concomitant risk of major bleeding (intracranial bleeding or fatal hemorrhages) make the use of this drugs problematic, with consequent underuse ⁵⁷. Older patients who take anticoagulants have an almost 2-fold increased risk of anticoagulation-related major and clinically relevant non-major bleeding than younger patients ⁹, possibly due to the higher prevalence of concomitant diseases, drug interactions, and age-related conditions. The multicenter, prospective Registro Informatizado de Enfermedad TromboEmbólica (RIETE) showed an increased rate of bleeding during anticoagulation therapy in patients aged ≥ 80 years old with acute VTE followed for at least three months after diagnosis compared to younger patients ⁵⁸. Vasco et al. ⁵⁹ described that in a subgroup of 610 people aged ≥ 90 years from the same RIETE registry, 4.9% had major bleeding complications, more than half fatal. Finally, the Worcester Venous Thromboembolism Study showed increased incidence rates of major bleedings in patients aged ≥ 65 years *versus* those < 65 years (9.2 and 4.8%, respectively) at 30 days after an acute episode of VTE. At 1 year, these proportions increased to 13.2 and 6.6% in older and younger people, respectively, with more than 50% of all major bleeding episodes occurring within 1 month of VTE diagnosis ⁷. Nevertheless, also the benefits in administering anticoagulants are increased in consideration of the high propensity of older subjects to hypercoagulability with a consequent higher risk of thrombosis. Thus, anticoagulants should be administered when indicated, such as prevention from stroke in older patients with atrial fibrillation (AF) or prevention of VTE expansion and recurrence in particular in older people with reduced mobility.

A further complication in using anticoagulation in older patients come from the fact that this population is underrepresented in many clinical trials of TE and of anticoagulant therapy ⁷ due to co-morbid conditions, short life expectancy, long-term immobility or contraindications to therapy ⁶⁰. As a consequence, treatment regimens derived from the results from clinical trials might not be optimal for older patients.

One of the most common indications to anticoagulants is atrial fibrillation (AF) in order to prevent stroke, since patients with AF have a 5-fold higher risk of ischemic stroke compared to healthy subjects, and the rate of thromboembolic events is substantially higher in the older people ⁶¹. An effective stroke prevention requires long-term use of oral anticoagulant with vitamin K antagonists (VKAs) or direct oral anticoagulants (DOACs)

and this evidence comes from randomized controlled trials (RCTs) demonstrating that warfarin reduces stroke/systemic embolism by 67%, ischemic stroke by 65%, and all-cause mortality by 26% compared with placebo or control⁶². This evidence has also been confirmed in the BAFTA trial, which specifically investigated older patients with AF ≥ 75 years, randomized to warfarin or aspirin 75 mg, and showed that warfarin significantly reduced thromboembolism by $> 50\%$, with no significant difference between warfarin and aspirin for major bleeding or intracranial bleeding⁶³. Nonetheless, few studies including very old subjects are available. Chao et al. investigated the risk of ischemic stroke and intracranial hemorrhage (ICH) and the net clinical benefit of oral anticoagulant treatment for very old patients with AF (≥ 90 years of age), demonstrating that antiplatelet agents were ineffective for stroke prevention and were not safer than no antithrombotic therapy; warfarin was associated with a positive net clinical benefit when compared to no antithrombotic therapy or antiplatelet drugs. Moreover, compared with warfarin, DOACs were associated with a lower risk of ICH, with no difference in risk of ischemic stroke⁶⁴. DOACs (apixaban, dabigatran, edoxaban and rivaroxaban) showed an improved risk-benefit profile in non-valvular AF patients as demonstrated by the lower incidence of stroke, intracranial hemorrhage, and death compared to warfarin⁶⁵. These agents are currently considered an alternative for VKAs to prevent stroke in patients with AF and have emerged as the preferred choice, particularly in patients newly started on anticoagulation⁶⁶. Additional evidence of efficacy and safety of DOACs came from a meta-analysis including 22 studies enrolling 440,281 AF patients ≥ 75 years that showed a reduction of the risk of stroke and systemic embolism compared to VKAs, although a comparable global incidence of major bleedings. In particular, DOACs reduced the rate of intracranial bleeding and hemorrhagic stroke events compared to VKAs, but increased the rate of GI bleedings, particularly rivaroxaban and dabigatran⁶⁷. Finally, a greater benefit of DOACs over VKA therapy has been found in subgroup analyses in subjects aged over 75 years, in patients with renal insufficiency (creatinine clearance 30-50 mL/min) and in those with a history of falls⁶⁸.

As far as VTE is concerned, all older patients with acute VTE and without any absolute contraindication to anticoagulation should receive active anticoagulant treatment as soon as diagnosis has been made, and even before in case of high clinical probability⁵⁸. This is important to prevent VTE expansion and recurrence. According to the new acute pulmonary embolism (PE) guidelines, the anticoagulant therapy should last no less than 3 months with the recommendation of an indefinite

duration of treatment for patients presenting with recurrent VTE⁶⁹. Once again, these indications are based on evidence coming from clinical trials not including older patients; thus, selection of the optimal antithrombotic agent, its dose, and duration of treatment is particularly difficult in this population. Anyway, during the last decade, DOACs have taken the place of VKAs in acute and long-term VTE treatment due to the convenience of administration and an excellent dose-response relationship without the need for monitoring or frequent dose adjustments⁷⁰. Indeed, the few reviews examining study evidence on DOACs compared to VKAs for VTE treatment in patients ≥ 75 years of age have demonstrated better efficacy and safety of DOACs over VKAs, with no increase in the risk of bleeding^{71,72}.

It must be noted that individual DOAC may have different profiles with respect to both effectiveness and safety. In the AMPLIFY study, apixaban (10 mg twice daily for seven days, then 5 mg twice daily for six months) was non inferior to warfarin bridged with enoxaparin in reduction of VTE in both the older and total patient population⁷³. The RE-COVER II trial found that dabigatran 150 mg twice daily was non inferior to warfarin, in reducing the risk of recurrent VTE in the older population⁷⁴. The Hokusai-VTE study, comparing edoxaban to warfarin, observed a significant reduction in risk of VTE compared with warfarin in the older patient population⁷⁵. In the EINSTEIN-DVT and PE trials rivaroxaban (15 mg twice daily for three weeks, then 20 mg once daily) was non inferior to warfarin bridged with enoxaparin in the reduction of VTE in both elderly populations and in the total patient population⁷⁶. Finally, in a meta-analysis of AF randomized controlled trials, DOACs reduced the risk of stroke or systemic thromboembolism by 19% compared with warfarin, largely due to a markedly lower rate of hemorrhagic stroke and intracranial bleeding (ICB)⁶⁵. As far as safety profiles concerned, it has been shown a significant reduction in risk of major bleeding for older patients taking apixaban or rivaroxaban compared to warfarin^{73,76}. In contrast, there was a non-significant difference in major bleeding for dabigatran and edoxaban compared with warfarin in the older and in the total patient population^{74,75}. In addition to these studies, very few real world studies have been conducted to compare the safety and effectiveness between DOACs focusing on very old patients. Deitelzweig et al. showed in a retrospective observational study of 88 582 very old patients (aged ≥ 80 y) with non-valvular AF that apixaban was associated with a lower risk of stroke/systemic embolism (SE), major bleeding (MB) and all cause mortality compared with dabigatran and rivaroxaban⁷⁷. A significant increase in the risk of gastrointestinal (GI) bleeding was observed in two randomized trials evaluating dabigatran and

rivaroxaban in AF patients: in the RE-LY trial the authors showed that dabigatran 150 mg bid (but not dabigatran 110 mg bid) was significantly associated with increased risk of major GI bleeding compared with warfarin ⁷⁸, whereas in the ROCKET-AF trial, rivaroxaban 20 mg once daily increased this annual risk by 1% ⁷⁹. DOAC-associated GI bleeding is probably related to the presence of the active drug in the GI tract thus facilitating bleeding from vulnerable lesions ⁸⁰. Thus, in patients with AF at a high-risk of GI bleeding, the 2016 European Society of Cardiology guidelines recommend using VKAs or DOACs other than dabigatran at a dose of 150 mg bid, rivaroxaban, 20 mg once daily, and edoxaban, 60 mg once daily ⁸¹.

An additional increased risk of bleeding in using DOACs is related to the possible presence of renal dysfunction, underweight or drugs interaction that are common among older subjects.

Currently, all DOACs have been approved for use in patients with chronic kidney disease (CKD) up to stage III. For patients with CrCl < 30 and > 15 ml/min, apixaban, rivaroxaban and edoxaban have been approved (with dose adjustment), while use of dabigatran is not recommended. In the European Union, none of the DOAC is indicated when CrCl is less than 15 ml/min or in patients undergoing hemodialysis, while for the latter group the FDA approved the use of apixaban and rivaroxaban (Tab. II) ⁸².

Being underweight has also been associated with an increased risk of major bleeding in patients taking oral anticoagulants, but there are still few evidences in this population, with consequent contrasting data. In fact, Park et al. showed, in an AF population taking DOACs according to their body mass index (BMI), an increased

risk of major bleeding and all-cause death in underweight patients, defined by BMI < 18.5, compared with being normal weight or obese. However, the median body weight in the low BMI group was 63.9 kg and the authors underline that a body weight < 60 kg was not an independent predictor of bleeding events ⁸³. More recently Lee et al. showed, in a large real-world Korean cohort with low body weight (< 60 kg) taking oral anticoagulants, that DOACs are better effectiveness and safety than warfarin both with regular and reduced dosage ⁸⁴. However, guidelines suggest caution with the use of non-vitamin K antagonist oral anticoagulants in patients with low body weight (< 60 kg) because of lack of evidences in this population (Tab. II).

Finally, polypharmacy, that is very frequent among older people, in addition to anticoagulant therapy may increase the risk of bleeding. Current knowledge of drug-drug interactions associated with DOACs mainly comes from animal studies, case reports, and limited pharmacokinetic measurement ⁸⁵. All the DOACs are substrates of P-glycoprotein (P-gp), and inhibition of this enzyme may increase bleeding risk. Thus, using a DOAC with P-gp inhibitors, such as ketoconazole and verapamil, or inducers, such a rifampin, may respectively increase or reduce the anticoagulant effect. In addition, rivaroxaban and apixaban also undergo metabolism to an extent of 40%-50% in the liver through the cytochrome P450 (CYP) pathway, specifically via CYP3A4, with a consequent possible interaction with drugs that induce or inhibit these pathways ⁸⁶ (Tab. II). VTE is of particular concern in patients admitted to hospital wards with acute medical conditions. In this population, patients aged > 75 years have an approximately 2-fold increased short-term risk of VTE than younger

Table II. DOACs special recommendation.

Anticoagulant	% Renal elimination	Drug interaction	Recommendation for dose reduction
Apixaban	25	CYP3A4; P-gp	2.5 mg BID if: • Age > 80 years • Weight < 60 kg • CrCl 15-30 ml/min • Not recommended if CrCl <15 ml/min
Rivaroxaban	67	CYP3A4; P-gp	15 mg QD if: • CrCl 15-50 ml/min • Not recommended if CrCl < 15 ml/min
Dabigatran	80	P-gp	75 mg QD if: • CrCl 30-50 ml/min • Not recommended if CrCl <30 ml/min • Age > 75 years
Edoxaban	35	P-gp	30 mg QD: • CrCl 15-50 ml/min • not recommended if CrCl <15 ml/min • Weight < 60 kg

patients⁹. Current guidelines recommend prophylaxis with fondaparinux or LMWH in high-risk hospitalized medical patients > 70 years to prevent VTE⁸⁷. Because age over 75 years is an independent risk factor of VTE in medically ill patients⁸⁸ the presence of only one additional acute medical condition, such as acute myocardial infarction, acute heart failure, active cancer requiring therapy, acute infectious disease, respiratory disease etc., is required to consider thromboprophylaxis for these patients⁸⁹. However, a recent case-control study conducted in France showed that in patients < 75 years there was a 2-fold increase in VTE in patients not receiving anticoagulants compared to those treated, while in patients > 75 years of age there were not differences in VTE risk in those taking or not taking anticoagulants⁹⁰. Thus, the authors agreed that thromboprophylaxis should only be prescribed following careful benefit-risk assessments, and that it is essential to consider major and non-major bleeding risks and comorbidities, including renal function, hypertension, infections and coronary artery disease.

In this context, The Padua Prediction Score (PPS) and the Improve Bleeding Score (IBS) have been approved by The American College of Chest Physicians 9th Edition guidelines as validated tools for VTE risk assessment in hospitalized old medical patients⁹¹. Some data suggest that a positive PPS and IBS are associated with early mortality in Internal Medicine patients⁹², but it has been demonstrated, in a group of old hospitalized patients, that their additional predictive accuracy is modest (positive PPS sensitivity: 96.67%, specificity: 20.74%; IBS sensitivity: 20.88%, specificity: 90.45%), resulting not very useful in clinical practice⁹³. Anyway, few studies about using these models in very old subjects are available.

CONCLUSIONS

This narrative review showed age-related changes of hemostasis pathways in older subjects, that can be considered as a normal phenomenon of aging, but may also be a consequence of dysregulated inflammatory pathways that contribute, together with multiple morbidities that are established prothrombotic risk factors, to increase the risk of thrombo-inflammatory disease in this population. Among the thrombotic cardiovascular diseases, stroke prevention remains one of the most important indication to use of anticoagulants even among very old and studies demonstrated the safety and efficacy of DOACs in terms of reduced risk of bleeding compared with warfarin. Anyway, despite the higher incidence, morbidity and mortality of TE in older subjects, a relevant proportion of this population do not receive prophylaxis with anticoagulants. In this context, a tailored therapy in older people

should be necessary taking into consideration changes attributed to aging and medical background in order to evaluate risk-benefit ratio in administering the potential optimal anticoagulant therapy.

References

- 1 Tracy RP, Bovill EG. Thrombosis and cardiovascular risk in the elderly. *Arch Pathol Lab Med* 1992;116:1307-12.
- 2 Benjamin EJ, Muntner P, Alonso A, et al. Heart disease and stroke statistics – 2019 update: a report from the American Heart Association. *Circulation* 2019;139. <https://doi.org/10.1161/CIR.0000000000000659>
- 3 Kung H-C, Hoyert DL, Xu J, et al. Deaths: final data for 2005. *Natl Vital Stat Rep* 2008;56:1-120.
- 4 Benjamin EJ, Blaha MJ, Chiuve SE, et al. Heart disease and stroke statistics – 2017 update: a report from the American Heart Association. *Circulation* 2017;135:e146-603. <https://doi.org/10.1161/CIR.0000000000000485>
- 5 Naess IA, Christiansen SC, Romundstad P, et al. Incidence and mortality of venous thrombosis: a population-based study. *J Thromb Haemost* 2007;5:692-9. <https://doi.org/10.1111/j.1538-7836.2007.02450.x>
- 6 Wilkerson WR, Sane DC. Aging and thrombosis. *Semin Thromb Hemost* 2002;28:555-68. <https://doi.org/10.1055/s-2002-36700>
- 7 Spencer FA, Gore JM, Lessard D, et al. Venous thromboembolism in the elderly. A community-based perspective. *Thromb Haemost* 2008;100:780-8.
- 8 Faller N, Limacher A, Méan M, et al. Predictors and causes of long-term mortality in elderly patients with acute venous thromboembolism: a Prospective Cohort study. *Am J Med* 2017;130:198-206. <https://doi.org/10.1016/j.amjmed.2016.09.008>
- 9 Tritschler T, Aujesky D. Venous thromboembolism in the elderly: a narrative review. *Thromb Res* 2017;155:140-147. <https://doi.org/10.1016/j.thromres.2017.05.015>
- 10 Cushman M. Epidemiology and risk factors for venous thrombosis. *Semin Hematol* 2007;44:62-9. <https://doi.org/10.1053/j.seminhematol.2007.02.004>
- 11 Gale AJ. Current understanding of hemostasis. *Toxicol Pathol* 2011;39:273-80. <https://doi.org/10.1177/0192623310389474>
- 12 Yamamoto K, Takeshita K, Kojima T, et al. Aging and plasminogen activator inhibitor-1 (PAI-1) regulation: implication in the pathogenesis of thrombotic disorders in the elderly. *Cardiovasc Res* 2005;66:276-85. <https://doi.org/10.1016/j.cardiores.2004.11.013>
- 13 Favaloro EJ, Franchini M, Lippi G. Aging hemostasis: changes to laboratory markers of hemostasis as we age – a narrative review. *Semin Thromb Hemost* 2014;40:621-33. <https://doi.org/10.1055/s-0034-1384631>
- 14 Tzoran I, Hoffman R, Monreal M. Hemostasis and thrombosis in the oldest old. *Semin Thromb Hemost* 2018;44:624-31. <https://doi.org/10.1055/s-0038-1657779>
- 15 Chung HY, Kim DH, Lee EK, et al. Redefining chronic

- inflammation in aging and age-related diseases: proposal of the senoinflammation concept. *Aging Dis* 2019;10:367-82. <https://doi.org/10.14336/AD.2018.0324>
- 16 Chung HY, Kim HJ, Kim KW, et al. Molecular inflammation hypothesis of aging based on the anti-aging mechanism of calorie restriction. *Microsc Res Tech* 2002;59:264-72. <https://doi.org/10.1002/jemt.10203>
 - 17 Chung HY, Lee EK, Choi YJ, et al. Molecular inflammation as an underlying mechanism of the aging process and age-related diseases. *J Dent Res* 2011;90:830-40. <https://doi.org/10.1177/0022034510387794>
 - 18 Chung HY, Sung B, Jung KJ, et al. The molecular inflammatory process in aging. *Antioxid Redox Signal* 2006;8:572-81. <https://doi.org/10.1089/ars.2006.8.572>
 - 19 Franceschi C, Campisi J. Chronic inflammation (inflammaging) and its potential contribution to age-associated diseases. *J Gerontol A Biol Sci Med Sci* 2014;69(Suppl 1):S4-9. <https://doi.org/10.1093/gerona/glu057>
 - 20 Baylis D, Bartlett DB, Patel HP, et al. Understanding how we age: insights into inflammaging. *Longev Healthspan* 2013;2:8. <https://doi.org/10.1186/2046-2395-2-8>
 - 21 Bauer ME, Fuente MD. The role of oxidative and inflammatory stress and persistent viral infections in immunosenescence. *Mech Ageing Dev* 2016;158:27-37. <https://doi.org/10.1016/j.mad.2016.01.001>
 - 22 Ershler WB. Interleukin-6: a cytokine for gerontologists. *J Am Geriatr Soc* 1993;41:176-81. <https://doi.org/10.1111/j.1532-5415.1993.tb02054.x>
 - 23 Oleksowicz L, Mrowiec Z, Zuckerman D, et al. Platelet activation induced by interleukin-6: evidence for a mechanism involving arachidonic acid metabolism. *Thromb Haemost* 1994;72:302-8.
 - 24 Mohebbi D, Kaplan D, Carlisle M, et al. Alterations in platelet function during aging: clinical correlations with thromboinflammatory disease in older adults. *J Am Geriatr Soc* 2014;62:529-35. <https://doi.org/10.1111/jgs.12700>
 - 25 Ziegler-Heitbrock L. The CD14+ CD16+ blood monocytes: their role in infection and inflammation. *J Leukoc Biol* 2007;81:584-92. <https://doi.org/10.1189/jlb.0806510>
 - 26 Seidler S, Zimmermann HW, Bartneck M, et al. Age-dependent alterations of monocyte subsets and monocyte-related chemokine pathways in healthy adults. *BMC Immunol* 2010;11:30. <https://doi.org/10.1186/1471-2172-11-30>
 - 27 Sadeghi HM, Schnelle JF, Thoma JK, et al. Phenotypic and functional characteristics of circulating monocytes of elderly persons. *Exp Gerontol* 1999;34:959-70. [https://doi.org/10.1016/s0531-5565\(99\)00065-0](https://doi.org/10.1016/s0531-5565(99)00065-0)
 - 28 Celermajer DS, Sorensen KE, Bull C, et al. Endothelium-dependent dilation in the systemic arteries of asymptomatic subjects relates to coronary risk factors and their interaction. *J Am Coll Cardiol* 1994;24:1468-74. [https://doi.org/10.1016/0735-1097\(94\)90141-4](https://doi.org/10.1016/0735-1097(94)90141-4)
 - 29 Chou TC, Yen MH, Li CY, et al. Alterations of nitric oxide synthase expression with aging and hypertension in rats. *Hypertension* 1998;31:643-8. <https://doi.org/10.1161/01.hyp.31.2.643>
 - 30 Taddei S, Virdis A, Ghiadoni L, et al. Age-related reduction of NO availability and oxidative stress in humans. *Hypertension* 2001;38:274-9. <https://doi.org/10.1161/01.hyp.38.2.274>
 - 31 Loscalzo J. Nitric oxide insufficiency, platelet activation, and arterial thrombosis. *Circ Res* 2001;88:756-62. <https://doi.org/10.1161/hh0801.089861>
 - 32 Garg UC, Hassid A. Nitric oxide-generating vasodilators and 8-bromo-cyclic guanosine monophosphate inhibit mitogenesis and proliferation of cultured rat vascular smooth muscle cells. *J Clin Invest* 1989;83:1774-7. <https://doi.org/10.1172/JCI114081>
 - 33 Balduini CL, Noris P. Platelet count and aging. *Haematologica* 2014;99:953-5. <https://doi.org/10.3324/haematol.2014.106260>
 - 34 Segal JB, Moliterno AR. Platelet counts differ by sex, ethnicity, and age in the United States. *Ann Epidemiol* 2006;16:123-30. <https://doi.org/10.1016/j.annepidem.2005.06.052>
 - 35 Le Blanc J, Lordkipanidzé M. Platelet function in aging. *Front Cardiovasc Med* 2019;6:109. <https://doi.org/10.3389/fcvm.2019.00109>
 - 36 Bastyr EJ, Kadrofske MM, Vinik AI. Platelet activity and phosphoinositide turnover increase with advancing age. *Am J Med* 1990;88:601-6. [https://doi.org/10.1016/0002-9343\(90\)90525-i](https://doi.org/10.1016/0002-9343(90)90525-i)
 - 37 Zahavi J, Jones NA, Leyton J, et al. Enhanced in vivo platelet "release reaction" in old healthy individuals. *Thromb Res* 1980;17:329-36. [https://doi.org/10.1016/0049-3848\(80\)90067-5](https://doi.org/10.1016/0049-3848(80)90067-5)
 - 38 Pastori D, Pignatelli P, Farcomeni A, et al. Aging-related decline of glutathione peroxidase 3 and risk of cardiovascular events in patients with atrial fibrillation. *J Am Heart Assoc* 2016;5. <https://doi.org/10.1161/JAHA.116.003682>
 - 39 Jin RC, Mahoney CE, Coleman Anderson L, et al. Glutathione peroxidase-3 deficiency promotes platelet-dependent thrombosis in vivo. *Circulation* 2011;123:1963-73. <https://doi.org/10.1161/CIRCULATIONAHA.110.000034>
 - 40 Zuo L, Prather ER, Stetskv M, et al. Inflammaging and oxidative stress in human diseases: from molecular mechanisms to novel treatments. *Int J Mol Sci* 2019;20. <https://doi.org/10.3390/ijms20184472>
 - 41 Favaloro EJ, Soltani S, McDonald J, et al. Cross-laboratory audit of normal reference ranges and assessment of ABO blood group, gender and age on detected levels of plasma coagulation factors. *Blood Coagul Fibrinolysis*.2005;16:597-605. <https://doi.org/10.1097/01.mbc.0000187250.32630.56>
 - 42 Kannel WB, Wolf PA, Castelli WP, et al. Fibrinogen and risk of cardiovascular disease. The Framingham Study. *JAMA* 1987;258:1183-6.
 - 43 Meade TW, North WR, Chakrabarti R, et al. Population-based distributions of haemostatic variables. *Br Med Bull* 1977;33:283-8. <https://doi.org/10.1093/oxfordjournals.bmb.a071448>
 - 44 Lowe GD, Rumley A, Woodward M, et al. Epidemiology of coagulation factors, inhibitors and activation markers:

- the Third Glasgow MONICA survey. I. Illustrative reference ranges by age, sex and hormone use. *Br J Haematol* 1997;97:775-84. <https://doi.org/10.1046/j.1365-2141.1997.1222936.x>
- 45 T K, Ad B, E B, Jp V, Fr R. Role of clotting factor VIII in effect of von willebrand factor on occurrence of deep-vein thrombosis. *Lancet* (London, England).
 - 46 Ofosu FA, Craven S, Dewar L, et al. Age-related changes in factor VII proteolysis in vivo. *Br J Haematol* 1996;94:407-12. <https://doi.org/10.1046/j.1365-2141.1996.d01-1793.x>
 - 47 Ten Cate H, Meade T. The Northwick Park Heart study: evidence from the laboratory. *J Thromb Haemost* 2014;12:587-92. <https://doi.org/10.1111/jth.12545>
 - 48 Tracy RP, Arnold AM, Ettinger W, et al. The relationship of fibrinogen and factors VII and VIII to incident cardiovascular disease and death in the elderly: results from the cardiovascular health study. *Arterioscler Thromb Vasc Biol* 1999;19:1776-83. <https://doi.org/10.1161/01.atv.19.7.1776>
 - 49 Kokame K, Sakata T, Kokubo Y, et al. von Willebrand factor-to-ADAMTS13 ratio increases with age in a Japanese population. *J Thromb Haemost* 2011;9:1426-8. <https://doi.org/10.1111/j.1538-7836.2011.04333.x>
 - 50 Favaloro EJ, Soltani S, McDonald J, et al. Laboratory identification of familial thrombophilia: do the pitfalls exceed the benefits? A reassessment of ABO-blood group, gender, age, and other laboratory parameters on the potential influence on a diagnosis of protein C, protein S, and antithrombin deficiency and the potential high risk of a false positive diagnosis. *Lab Hematol* 2005;11:174-84. <https://doi.org/10.1532/LH96.05029>
 - 51 Bauer KA, Weiss LM, Sparrow D, et al. Aging-associated changes in indices of thrombin generation and protein C activation in humans. Normative Aging Study. *J Clin Invest* 1987;80:1527-34. <https://doi.org/10.1172/JCI113238>
 - 52 Hamilton PJ, Dawson AA, Ogston D, et al. The effect of age on the fibrinolytic enzyme system. *J Clin Pathol* 1974;27:326-9. <https://doi.org/10.1136/jcp.27.4.326>
 - 53 Dolan G, Neal K, Cooper P, et al. Protein C, antithrombin III and plasminogen: effect of age, sex and blood group. *Br J Haematol* 1994;86:798-803. <https://doi.org/10.1111/j.1365-2141.1994.tb04832.x>
 - 54 Yamamoto K, Takeshita K, Shimokawa T, et al. Plasminogen activator inhibitor-1 is a major stress-regulated gene: implications for stress-induced thrombosis in aged individuals. *Proc Natl Acad Sci USA* 2002;99:890-5. <https://doi.org/10.1073/pnas.022608799>
 - 55 Browder T, Folkman J, Pirie-Shepherd S. The hemostatic system as a regulator of angiogenesis. *J Biol Chem* 2000;275:1521-4. <https://doi.org/10.1074/jbc.275.3.1521>
 - 56 Folkman J, Browder T, Palmblad J. Angiogenesis research: guidelines for translation to clinical application. *Thromb Haemost* 2001;86:23-33.
 - 57 Hylek EM, Evans-Molina C, Shea C, et al. Major hemorrhage and tolerability of warfarin in the first year of therapy among elderly patients with atrial fibrillation. *Circulation* 2007;115:2689-96. <https://doi.org/10.1161/CIRCULATIONAHA.106.653048>
 - 58 Palareti G, Poli D. The prevention of venous thromboembolism recurrence in the elderly: a still open issue. *Expert Rev Hematol* 2018;11:903-9. <https://doi.org/10.1080/17474086.2018.1526667>
 - 59 Vasco B, Villalba JC, Lopez-Jimenez L, et al. Venous thromboembolism in nonagenarians. Findings from the RIETE Registry. *Thromb Haemost* 2009;101:1112-8.
 - 60 Lacruz B, Tiberio G, Núñez MJ, et al. Venous thromboembolism in centenarians: Findings from the RIETE registry. *Eur J Intern Med* 2016;36:62-6. <https://doi.org/10.1016/j.ejim.2016.07.025>
 - 61 Lip GYH, Nieuwlaat R, Pisters R, et al. Refining clinical risk stratification for predicting stroke and thromboembolism in atrial fibrillation using a novel risk factor-based approach: the euro heart survey on atrial fibrillation. *Chest* 2010;137:263-72. <https://doi.org/10.1378/chest.09-1584>
 - 62 Hart RG, Pearce LA, Aguilar MI. Meta-analysis: antithrombotic therapy to prevent stroke in patients who have non-valvular atrial fibrillation. *Ann Intern Med* 2007;146:857-67. <https://doi.org/10.7326/0003-4819-146-12-200706190-00007>
 - 63 Mant J, Hobbs FDR, Fletcher K, et al. Warfarin versus aspirin for stroke prevention in an elderly community population with atrial fibrillation (the Birmingham Atrial Fibrillation Treatment of the Aged Study, BAFTA): a randomised controlled trial. *Lancet* 2007;370:493-503. [https://doi.org/10.1016/S0140-6736\(07\)61233-1](https://doi.org/10.1016/S0140-6736(07)61233-1)
 - 64 Chao T-F, Liu C-J, Lin Y-J, et al. Oral anticoagulation in very elderly patients with atrial fibrillation: a Nationwide Cohort study. *Circulation* 2018;138:37-47. <https://doi.org/10.1161/CIRCULATIONAHA.117.031658>
 - 65 Ruff CT, Giugliano RP, Braunwald E, et al. Comparison of the efficacy and safety of new oral anticoagulants with warfarin in patients with atrial fibrillation: a meta-analysis of randomised trials. *Lancet* 2014;383(9921):955-962. [doi:10.1016/S0140-6736\(13\)62343-0](https://doi.org/10.1016/S0140-6736(13)62343-0)
 - 66 Steffel J, Verhamme P, Potpara TS, et al. The 2018 European Heart Rhythm Association Practical Guide on the use of non-vitamin K antagonist oral anticoagulants in patients with atrial fibrillation. *Eur Heart J* 2018;39:1330-93. <https://doi.org/10.1093/eurheartj/ehy136>
 - 67 Silverio A, Di Maio M, Protta C, et al. Safety and efficacy of non-vitamin K antagonist oral anticoagulants in elderly patients with atrial fibrillation. *Eur Heart J Cardiovasc Pharmacother*. December 2019. <https://doi.org/10.1093/ehjcvp/pvz073>
 - 68 Hanon O, Jeandel C, Jouanny P, et al. Anticoagulant treatment in elderly patients with atrial fibrillation: position paper. *Geriatr Psychol Neuropsychiatr Vieil* 2019;17:341-55. <https://doi.org/10.1684/pnv.2019.0834>
 - 69 Konstantinides SV, Meyer G, Becattini C, et al. 2019 ESC Guidelines for the diagnosis and management of acute pulmonary embolism developed in collaboration with the European Respiratory Society (ERS). *Eur Heart J* 2020;41:543-603. <https://doi.org/10.1093/eurheartj/ehz405>

- 70 Kearon C, Akl EA, Ornelas J, et al. Antithrombotic therapy for VTE disease: CHEST Guideline and Expert Panel Report. *Chest* 2016;149:315-52. <https://doi.org/10.1016/j.chest.2015.11.026>
- 71 Geldhof V, Vandenbriele C, Verhamme P, et al. Venous thromboembolism in the elderly: efficacy and safety of non-VKA oral anticoagulants. *Thromb J* 2014;12:21. <https://doi.org/10.1186/1477-9560-12-21>
- 72 Sadlon AH, Tsakiris DA. Direct oral anticoagulants in the elderly: systematic review and meta-analysis of evidence, current and future directions. *Swiss Med Wkly* 2016;146:w14356. <https://doi.org/10.4414/smww.2016.14356>
- 73 Agnelli G, Buller HR, Cohen A, et al. Apixaban for extended treatment of venous thromboembolism. *N Engl J Med* 2013;368:699-708. <https://doi.org/10.1056/NEJMoa1207541>
- 74 Schulman S, Kakkar AK, Goldhaber SZ, et al. Treatment of acute venous thromboembolism with dabigatran or warfarin and pooled analysis. *Circulation* 2014;129:764-72. <https://doi.org/10.1161/CIRCULATIONAHA.113.004450>
- 75 Vanassche T, Verhamme P, Wells PS, et al. Impact of age, comorbidity, and polypharmacy on the efficacy and safety of edoxaban for the treatment of venous thromboembolism: an analysis of the randomized, double-blind Hokusai-VTE trial. *Thromb Res* 2018;162:7-14. <https://doi.org/10.1016/j.thromres.2017.12.005>
- 76 Prins MH, Lensing AW, Bauersachs R, et al. Oral rivaroxaban versus standard therapy for the treatment of symptomatic venous thromboembolism: a pooled analysis of the EINSTEIN-DVT and PE randomized studies. *Thromb J* 2013;11:21. <https://doi.org/10.1186/1477-9560-11-21>
- 77 Deitelzweig S, Keshishian A, Li X, et al. Comparisons between oral anticoagulants among older nonvalvular atrial fibrillation patients. *J Am Geriatr Soc* 2019;67:1662-71. <https://doi.org/10.1111/jgs.15956>
- 78 Eikelboom JW, Wallentin L, Connolly SJ, et al. Risk of bleeding with 2 doses of dabigatran compared with warfarin in older and younger patients with atrial fibrillation: an analysis of the randomized evaluation of long-term anticoagulant therapy (RE-LY) trial. *Circulation* 2011;123:2363-72. <https://doi.org/10.1161/CIRCULATIONAHA.110.004747>
- 79 Patel MR, Mahaffey KW, Garg J, et al. Rivaroxaban versus warfarin in nonvalvular atrial fibrillation. *N Engl J Med* 2011;365:883-91. <https://doi.org/10.1056/NEJMoa1009638>
- 80 Heidbuchel H, Verhamme P, Alings M, et al. Updated European Heart Rhythm Association Practical Guide on the use of non-vitamin K antagonist anticoagulants in patients with non-valvular atrial fibrillation. *Europace* 2015;17:1467-507. <https://doi.org/10.1093/europace/euv309>
- 81 Kirchhof P, Benussi S, Kotecha D, et al. 2016 ESC Guidelines for the management of atrial fibrillation developed in collaboration with EACTS. *Eur Heart J* 2016;37:2893-62. <https://doi.org/10.1093/eurheartj/ehw210>
- 82 Weber J, Olyaei A, Shatzel J. The efficacy and safety of direct oral anticoagulants in patients with chronic renal insufficiency: a review of the literature. *Eur J Haematol* 2019;102:312-8. <https://doi.org/10.1111/ejh.13208>
- 83 Park CS, Choi E-K, Kim HM, et al. Increased risk of major bleeding in underweight patients with atrial fibrillation who were prescribed non-vitamin K antagonist oral anticoagulants. *Heart Rhythm* 2017;14:501-7. <https://doi.org/10.1016/j.hrthm.2016.12.036>
- 84 Lee S-R, Choi E-K, Park CS, et al. Direct oral anticoagulants in patients with nonvalvular atrial fibrillation and low body weight. *J Am Coll Cardiol* 2019;73:919-31. <https://doi.org/10.1016/j.jacc.2018.11.051>
- 85 Chang S-H, Chou I-J, Yeh Y-H, et al. Association between use of non-vitamin K oral anticoagulants with and without concurrent medications and risk of major bleeding in non-valvular atrial fibrillation. *JAMA* 2017;318:1250-9. <https://doi.org/10.1001/jama.2017.13883>
- 86 Bethishou L, Gregorian T, Won K, et al. Management of venous thromboembolism in the elderly: a review of the non-vitamin K oral anticoagulants. *Consult Pharm* 2018;33:248-61. <https://doi.org/10.4140/TCP.n.2018.248>
- 87 Di Nisio M, Porreca E. Prevention of venous thromboembolism in hospitalized acutely ill medical patients: focus on the clinical utility of (low-dose) fondaparinux. *Drug Des Devel Ther* 2013;7:973-80. <https://doi.org/10.2147/DDDT.S38042>
- 88 Alikhan R, Cohen AT, Combe S, et al. Risk factors for venous thromboembolism in hospitalized patients with acute medical illness: analysis of the MEDENOX study. *Arch Intern Med* 2004;164:963-8. <https://doi.org/10.1001/archinte.164.9.963>
- 89 Lacut K, Le Gal G, Mottier D. Primary prevention of venous thromboembolism in elderly medical patients. *Clin Interv Aging* 2008;3:399-411. <https://doi.org/10.2147/cia.s832>
- 90 Hemon F, Fouchard F, Tromeur C, et al. Association between hospitalization for acute medical illness and VTE risk: a lower efficacy of thromboprophylaxis in elderly patients? Results from the EDITH case-control study. *Eur J Intern Med* 2017;44:39-43. <https://doi.org/10.1016/j.ejim.2017.05.029>
- 91 Depietri L, Marietta M, Scarlini S, et al. Clinical impact of application of risk assessment models (Padua Prediction Score and Improve Bleeding Score) on venous thromboembolism, major hemorrhage and health expenditure associated with pharmacologic VTE prophylaxis: a "real life" prospective and retrospective observational study on patients hospitalized in a Single Internal Medicine Unit (the STIME study). *Intern Emerg Med* 2018;13:527-34. <https://doi.org/10.1007/s11739-018-1808-z>
- 92 La Regina M, Orlandini F, Marchini F, et al. Combined assessment of thrombotic and haemorrhagic risk in acute medical patients. *Thromb Haemost* 2016;115:392-8. <https://doi.org/10.1160/TH14-12-1050>
- 93 Arpaia GG, Caleffi A, Marano G, et al. Padua prediction score and IMPROVE score do predict in-hospital mortality in Internal Medicine patients. *Intern Emerg Med*; Jan 2020. <https://doi.org/10.1007/s11739-019-02264-4>