

REVIEW

Mechanisms and potential treatments for declining olfactory function and neurogenesis in the ageing brain

K.D. Broad

UCL Institute of Ophthalmology, University College, London, UK

The role of olfactory function in maintaining quality of life and as a potential surrogate marker of neurogenic activity in the elderly brain is an underappreciated topic. The olfactory system is complex and is unusual in that its function is maintained by neurogenesis at multiple sites throughout the lifetime of an organism, which in humans may be over 80 years in length. Declines in olfactory function are common with advancing age and this is associated with reductions in the quality of life, the perception of flavour and neurogenesis. These reductions in neurogenesis may simply be a consequence of advancing age or may reflect the nascent development of underlying neurological dysfunction. There are a number of potential therapeutic interventions that can result in a gain of olfactory function, these range from behavioural modification, to dietary supplementation and pharmaceutical intervention but they are all thought to work in part by increasing hippocampal and olfactory neurogenesis. This review discusses the mechanisms underlying olfactory decline in the elderly, reviews a number of potential strategies for improving olfactory function and hypotheses that increasing the rate of neurogenesis in the ageing brain would improve the quality of life not only by improving olfaction but by improving a range of cognitive processes that are dependent on neurogenesis including mood.

Key words: Olfaction, Ageing, Neurogenesis, Depression, Flavour

INTRODUCTION

A declining ability to detect, identify and correctly interpret the environmental significance of olfactory cues is a common feature of advancing age¹⁻⁹. This often leads to a decline in the quality of life, reduced environmental awareness and a reduction in the appreciation of “taste” and “flavour”, which is thought to result in a diminished appetite and nutritional neglect. In elderly human populations attempts to definitively link olfactory decline with nutritional neglect have yielded ambiguous results¹⁰⁻¹³. This is despite the fact that a high incidence of “taste” dysfunction is actually reflective of declines in olfactory function¹⁴. And that the perception of “flavour” is thought to be generated by a multimodal integration of taste, olfactory, textural and temperature information in a

distributed network encompassing the tractus nucleus solitarius, insula, operculum, orbitofrontal, pyriform, and anterior cingulate cortices^{15,16}. Despite a degree of ambiguity concerning the exact relationship between olfactory decline and nutritional neglect a number of studies have shown that declines in olfactory function are associated with an increased risk of mortality in the elderly¹⁷⁻²⁰. One recent study in particular demonstrated that declines in olfactory function were strongly associated with an increased risk of mortality but declines in visual or auditory function were not¹⁷. It has long been presumed that a decline in olfactory function is part of the normal ageing process¹⁻¹⁴ that parallels a generalised decline in sensory and cognitive function that occurs in the absence of obvious neurological disease states and impaired olfactory function does predict generalised cognitive decline in the elderly²¹. However, a number of diseases common in the

■ Received: May 11, 2016 - Accepted: July 22, 2016

■ Correspondence: Kevin Broad, UCL Institute of Ophthalmology, University College London, 11-43 Bath Street, London EC1V 9EL, UK - E-mail: kev.broad@ucl.ac.uk

elderly, are also associated with olfactory dysfunction. These include depression²²⁻²⁵, Alzheimer's disease²⁴⁻²⁷, prion disease²⁶, epilepsy²⁸⁻²⁹, Parkinson's disease³⁰⁻³¹, stroke³²⁻³³, physical trauma³⁴ and diabetes³⁵⁻³⁶. In some instances olfactory decline precedes the onset of clinical symptoms by a considerable margin³⁰⁻³¹, suggesting that the olfactory declines seen in "normal" ageing patients may also be contributed to by the nascent development of neurological diseases often seen in the elderly.

The obvious questions that arise from these observations are: What mechanisms contribute to a decline in olfactory function in the "normal" ageing brain? Are there practical strategies for improving olfactory function? And would improving olfactory function lead to an increase in the quality of life and improvements in health in the elderly population? In this review we focus on the importance of a number of mechanisms that contribute to olfactory decline and examine the development of a number of strategies that may lead to a gain in olfactory function and subsequent improvements in the quality of life.

CHANGES IN THE OLFACTORY SYSTEM DURING THE AGEING PROCESS

An appreciation of how the human olfactory system extracts physiologically and behaviourally salient information from the environment is complicated by the fact that most research into olfactory function is conducted in rodents who possess two functional and neuroanatomically distinct olfactory systems. The vomeronasal system in rodents is thought to be primarily responsible for the transmission of inter-species reproductive, social and physiological salient information transmitted by pheromones contained in urine. The alternative main olfactory system is considered to function as an analyser of "general" volatile odours. In primates, the vomeronasal system becomes evolutionarily redundant around the time that trichromatic colour vision evolves³⁷ and as a consequence the human olfactory system relies on the main olfactory system only. This can make extrapolation of data derived from rodents problematic, but the main olfactory and vomeronasal epithelial systems in rodents are functionally very similar in that their receptor neurons are generated from stem cells throughout the life of the organism and differentiate into neurons that express 1-2 single olfactory receptors per cell.

THE OLFACTORY EPITHELIUM

In humans the main olfactory epithelium is localised in the nasal cavity and is positioned around the cribriform plate. Its olfactory receptor neurons are unusual

in that they continue to be generated throughout life from stem/progenitor cells, which differentiate into odorant-detecting olfactory receptor neurons which send projections into the mucous layer to sample the olfactory environment and unmyelinated axons through the cribriform plate to transfer information to the olfactory bulb³⁸⁻³⁹. On differentiation, each olfactory receptor neuron expresses one olfactory selected from around 1,000 subtly different G-protein coupled odorant receptor genes³⁹⁻⁴¹. The proximal cues that influence this selection process are poorly understood but in the functionally similar but closely studied murine vomeronasal system the differentiation and selection process the cue is urinary pheromones, which act to initiate differentiation and influence the receptor selection process⁴²⁻⁴³. During ageing a number of changes are thought to occur to the olfactory epithelium primarily the number of differentiating stem cells are reduced⁴⁴⁻⁴⁷, a process that in rodents occurs in parallel with reductions in the differentiation of vomeronasal stem cells⁴⁸⁻⁵⁰. This decline in neural differentiation is accompanied with a decline in the expression of *Asc11*, a pro-neural gene required for generation of olfactory sensory neurons⁵¹ and epidermal growth factor signalling³⁹. The olfactory epithelium also demonstrates remarkable anatomical⁵² and functional⁵³ regenerative capabilities. Following olfactory lesions anatomical recovery is evident with ~45 days and odorant expression patterns are re-established within 90 days⁵⁴. It has been hypothesised that there may be a reduction in the regenerative capacity of the olfactory epithelium as an animal ages and this is supported by evidence in the mouse. The efficacy of the regenerative process following a methimazole-induced lesion at the age of 16 months was considerably lower than that occurring at 10 days or 3 months old, the number of olfactory receptor neurons being around a third of those in the younger groups⁵⁵.

Two potential consequences of age that these studies fail to address is that the epigenetic stimuli that induce the transformation of olfactory stem cells into functional receptor neurons are unclear, in the mouse vomeronasal system this stimulus is urinary pheromones⁴²⁻⁴³. The mouse vomeronasal organ also expresses 44 main olfactory receptors with the same 1 olfactory receptor per neuron expression pattern displayed in the main olfactory epithelium and the differentiation and olfactory receptor expression of these neurons is also under the control of urinary pheromones⁴³. In both systems differentiation is reduced with age, in the vomeronasal system this may reflect a reduction in the sensitivity of vomeronasal stem cells to the transformative epigenetic effects of urinary pheromones, and this reduction in sensitivity to the environmental epigenetic cues that influence differentiation may also occur in the main olfactory

epithelium. Delivering drugs targeted to the olfactory epithelium by an aerosol is technically very feasible and investigation of the epigenetic mechanisms that control the differentiation of olfactory stem cells into functional olfactory receptor neurons may lead to treatments that support increased olfactory function in the elderly. One other problem that these studies do not address is that olfactory receptor neurons send unmyelinated projection to the glomerular layer of the olfactory bulb, this is a difficult process requiring successful axonal targeting and transport over a considerable distance. Although this is an underexplored topic in the olfactory system, these axonal processes in other models decline considerably with age⁵⁶⁻⁵⁸ and it is likely that a similar process occurs in the olfactory system.

PRIMARY OLFACTORY PROCESSING. THE OLFACTORY BULB

The main olfactory system for an underappreciated sensory modality is surprisingly complex and many of the important but subjective experiences with an olfactory component including the perception of “palatability” or “flavour” rely on multisensory integration of olfactory with other inputs including taste, texture and temperature¹⁶, which adds a further layer of complexity to attempts to appreciate olfactory function. A diagram illustrating the basic structure of the main olfactory system is outlined in Figure 1.

Axonal processes from olfactory receptor neurons synapse on the dendrites of the primary projection neurons of the olfactory bulb, the mitral and tufted cells, which together form a structure called the glomerulus. They are excitatory glutaminergic neurons, their dendrites project to a single glomerulus, and as well as receiving input from olfactory receptor neurons they have extensive input from lateral dendrites. Despite being excitatory neurons, the modification of these cells occurs by reciprocal dendrodendritic connections with γ -aminobutyric acid (GABA)ergic granule cells, the

activity of which is modulated by centrifugal input from neurons outside the olfactory bulb. The incoming axons from the olfactory receptor neurons also synapse on local GABAergic interneurons (periglomerular cells) that are activated by glutamate released from mitral and tufted cells and provide an inhibitory influence within the glomerulus. Glomeruli are the first synaptic relay in the olfactory pathway and play a basic role in smell perception. A second level of olfactory processing occurs at the granular layer of the olfactory bulb by inhibitory GABAergic neurons that are activated by glutamate released from lateral dendrites of mitral cells, which act to inhibit the network in contrast to enhancement between mitral cells⁵⁹. The olfactory bulb is also unusual in that its neurons are replenished throughout life by neuroblasts generated in the sub ventricular zone and transported via the rostral migratory stream⁶⁰.

The volume of the olfactory bulb and the number of its laminae are thought to decline with age, but this decline in volume is thought to interact with a number of environmental influences including smoking⁶¹⁻⁶³. Olfactory bulb volume is a gross measurement but correlations have been found between volume and odour recognition thresholds in the normal ageing brain⁶². Changes in the expression of proteins normally thought to be associated with dementia may also increase in the “normal” ageing brain. The proliferation of neurodegenerative fibrillary tangles occurs in 35.5 to 40.5% of non-demented older patients⁶⁴ and the expression of α -synuclein increases as a consequence of ageing in the marmoset brain⁶⁵. The efficient function of the olfactory bulb is also dependent on the continuous replenishment of neurons generated in the sub-ventricular zone, a process that is reduced in number and disrupted with regard to spatiotemporal organisation as a consequence of age^{66 67}.




Figure 1. Simplified structure of the main olfactory system, showing basic interconnectivity and convergence on the frontal cortex.

SECONDARY AND TERTIARY OLFACTORY PROCESSING.

A DISTRIBUTED NETWORK

Axons from the olfactory bulb project into a distributed olfactory processing network. The engagement of each part of this network contributes to the processing of olfactory cues, the formation and recall of olfactory memories, the exact contributions of each stage of this network are beyond the scope of this review and are extensively covered elsewhere⁶⁸. The first relays in this network are the anterior olfactory nucleus, olfactory tubercle and the pyriform cortex, (a large planar cortical olfactory area)⁶⁹. The functional interconnectivity of the network is complex but olfactory information from these three areas are then routed through a distributed network encompassing the amygdala⁷⁰⁻⁷², entorhinal cortex^{73 74}, hippocampus⁷⁵ and the frontal cortex^{76 77}. Many of these areas send projections back to the olfactory bulb where they terminate primarily in the granule layer.

A number of changes occur within this network as a consequence of ageing, these include a reduction in the volume of the hippocampus, amygdala, pyriform cortex and accessory olfactory nucleus⁷⁸. The volume, neuronal organisation and localisation of the Islands of Calleja is altered by advancing age and is thought to contribute to a decline in olfactory function⁷⁹. Neurofibrillary and α -synuclein pathology in the olfactory bulb is associated with olfactory dysfunction in elderly non-demented patients, which is likely to compromise the ability of mitral and tufted cells to project to their appropriate targets^{64 65}. The olfactory system with its dependence on lifelong neural replenishment and the requirement that these new neurons be correctly integrated into existing neural structures is a classic example of extreme neuroplasticity. The mechanisms underlying neuroplasticity are extremely complex and warrant a review of their own but the mechanisms mediating neuroplasticity do decline with age. The expression of brain derived neurotrophic factor also declines with age⁸⁰ which is likely to compromise the ability of new neurons to integrate successfully into pre-existing networks.

THE AGEING OLFACTORY SYSTEM AS A THERAPEUTIC TARGET

The olfactory system is unusual in that its function is dependent on the continuous replenishment of neurons throughout the lifetime of an individual. In elderly humans, this process therefore needs to be maintained over an 80 year period, which given the relative decline in other cognitive processes is an impressive feat. This occurs in four levels of the olfactory system:

the olfactory epithelium³⁷⁻⁴³, the olfactory bulb⁸¹, sub ventricular zone, which provides replacement GABA-nergic interneurons to the olfactory bulb^{60 66 67} and the dentate gyrus of the hippocampus^{82 83}. Increasing neurogenesis does provide a route to improve olfactory function in the elderly, which also has the advantage that it may improve other indices of life quality, including mood. The importance of olfaction to human wellbeing is underappreciated but some interventions that may improve olfactory function are simple and may provide other benefits.

Behavioural modification; It is well recognised that increases in physical activity promote increased neurogenesis and immature neuron survival in both the hippocampus⁸⁴ and sub-ventricular system olfactory system⁸⁵ but not in the olfactory bulb itself⁸⁶. A number of different mechanisms have been proposed to mediate the interactions between exercise and adult neurogenesis including brain derived neurotrophic factor^{87 88}, insulin-like growth factor 1 (IGF1)^{89 90}, vascular endothelial growth factor (VEGF)⁹¹ and the Wnt signalling pathway^{92 93}. Neurogenesis induced by exercise is also thought to function as one of the experience driven mechanisms that promote neural plasticity⁹⁴. For obvious reasons conclusive attempts to link exercise, neurogenesis and increases in olfactory function in elderly patients are problematic but human epidemiological based studies have demonstrated that exercise does reduce olfactory decline in elderly patients⁹⁵. The olfactory system expressing high levels of plasticity is responsive to its environment. Olfactory environmental enrichment is also associated with increases in neurogenesis^{96 97}, improvements in short-term odour memory⁹⁸ and olfactory discrimination^{99 100}. This work has been undertaken in rodents and the practical extrapolation of this approach to elderly humans is unclear. Olfactory function is also decreased in the obese, a phenotype that is likely to interact and be associated with ageing and low activity levels^{101 102}. Hippocampal neurogenesis is thought to be reduced in depression which may contribute to the development of deficits in olfactory function²²⁻²⁵, one effect of antidepressants is that they increase neurogenesis and this is probably contributed too by reductions in stress that occur with an increase in mood. Treating depression in the elderly by alternative non-pharmaceutical means including cognitive behavioural therapies may also contribute to improvements in olfactory function¹⁰³⁻¹⁰⁵.

Pharmacological treatment; a range of pharmacological interventions may induce gains in olfactory function, these range from dietary supplementation to more traditional pharmacological treatments.

The consumption of omega-3 fatty acids increases olfactory function^{106 107}, an effect that may be associated

with increased neurogenesis¹⁰⁸. A range of other dietary supplements may also promote neurogenesis which as an advantageous side effect may improve olfactory function, these include curcumin¹⁰⁹, flavonoids^{110 111}, vitamin A¹¹² and caffeine¹¹³.

Olfactory function and neurogenesis is also reduced in depression and it is noteworthy that anti-depressives are thought to work in part by increasing neurogenesis in the dentate gyrus^{22-25 103-105}. However it is thought that they also induce increases in olfactory neurogenesis and olfactory function in rodents¹⁰⁴, this is likely to be an effect replicated in humans as recovery from depression is thought to result in gains of olfactory function¹⁰⁵. Other drugs can also induce olfactory neurogenesis and differentiation, these include valproic acid and other histone deacetylase inhibitors used to treat epilepsy^{114 115}. Two other non-pharmacological interventions for depression also increase neurogenesis and increase olfactory function, electroconvulsive therapy and transcranial magnetic stimulation¹¹⁶⁻¹¹⁹.

CONCLUSIONS

The neural systems supporting olfactory function are complex and are unusual in that they rely on the continuous neurogenesis throughout life. This occurs at four separate sites, the olfactory epithelium³⁸⁻⁴³, endogenously in the olfactory bulb⁸¹, the sub-ventricular zone^{60 66 67} and the dentate gyrus^{82 83}. Olfactory function declines with advancing age and this is thought to reflect a reduction in this process^{1-14 44-47}. A range of neurological disorders common in the elderly are also associated with reductions in neurogenesis and a decline in olfactory function may therefore be a surrogate marker or a harbinger of a developing neurological condition²²⁻³⁵. Olfaction is an underappreciated sensory modality in humans but declines in olfactory function are predictive of mortality in the elderly, whilst declines in the more appreciated senses of vision and auditory function are not¹⁷⁻²¹. This may reflect the reliance of olfaction on neurogenesis, a decline in which may be a robust surrogate marker of declining neurological function. Declines in olfactory function also lead to a reduction in the quality of life including the reduction of “flavour” or “palatability”, which in turn may be reflective of this decline in neurogenesis¹⁰⁻¹⁶. There are a number of potential mechanisms by which olfactory function may be improved in the elderly and given olfaction’s reliance on neurogenesis it is unsurprising that many of these are thought to work by improving this function. Olfaction is an underappreciated sensory modality and its role in the maintenance of a high quality of life and as a potential harbinger of mortality in the elderly is an

underexplored topic that warrants further investigation. In particular improvement of neurogenesis in the elderly by a variety of approaches may yield improvements in a range of cognitive functions in parallel with olfaction including mood.

References

- Murphy C. *The chemical senses and nutrition in older adults*. J Nutr Elder 2008;27:247-65.
- Doty RL. *The olfactory system and its disorders*. Semin Neurol 2009;29:74-81.
- Rawson NE, Gomez G, Cowart BJ, et al. *Age-associated loss of selectivity in human olfactory sensory neurons*. Neurobiol Aging 2012;33:1913-9.
- Brämerson A, Johansson L, Ek L, et al. *Prevalence of olfactory dysfunction: the Skövde population based study*. Laryngoscope 2004;114:733-7.
- Hüttenbrink KB, Hummel T, Berg D, et al. *Olfactory dysfunction: common in later life and early warning of neurodegenerative disease*. Dtsch Arztebl Int 2013;110:1-7.
- Karpa MJ, Gopinath B, Rochtchina E, et al. *Prevalence and neurodegenerative or other associations with olfactory impairment in an older community*. J Aging Health 2010;22:154-68.
- Schubert CR, Cruickshanks KJ, Fischer ME, et al. *Odor identification and cognitive function in the Beaver Dam Offspring Study*. J Clin Exp Neuropsychol 2013;35:669-76.
- Doty RL, Kamath V. *The influences of age on olfaction: a review*. Front Psychol 2014;5:20.
- Lafreniere D, Mann N. *Anosmia: loss of smell in the elderly*. Otolaryngol Clin North Am 2009;42:123-31.
- Ferris AM, Duffy VB. *Effects of olfactory deficits on nutritional status. Does age predict persons at risk?* Ann NY Acad Sci 1989;561:113-23.
- Duffy VB, Backstrand JR, Ferris AM. *Olfactory dysfunction and related nutritional risk in free-living, elderly women*. J Am Diet Assoc 1995;95:879-84.
- Mattes RD. *The chemical senses and nutrition in aging: challenging old assumptions*. J Am Diet Assoc 2002;102:192-6.
- Toussaint N, de Roon M, van Campen JPC, et al. *Loss of olfactory function and nutritional status in vital older adults and geriatric patients*. Chem Senses 2015;40:197-203.
- Deems DA, Doty RL, Settle RG, et al. *Smell and taste disorders, a study of 750 patients from the University of Pennsylvania Smell and Taste Centre*. Arch Otolaryngol Head Neck Surg 1991;117:519-28. <http://www.ncbi.nlm.nih.gov/pubmed/2021470>
- Escanilla O, Victor JD, Di Lorenzo PM. *Odor-taste convergence in the nucleus of the solitary tract of the awake freely licking rat*. J Neurosci 2015;35:6284-97.
- Small DM, Green BG. *A proposed model of a flavour modality*. In: Murray MM, Wallace MT, Eds. *The neural basis of multisensory processes*. Boca Raton, FL: CRC 2012. Chap 36.

- 17 Schubert CR, Fischer ME, Pinto AA, et al. *Sensory impairments and the risk of mortality in older adults*. J Gerontol A Biol Sci Med Sci 2017;72:710-5.
- 18 Devenand DP, Lee S, Manly J, et al. *Olfactory identification deficits and increased mortality in the community*. Ann Neurol 2015;78:401-11.
- 19 Pinto JM, Wroblewski KE, Kern DW, et al. *Olfactory dysfunction predicts 5-year mortality in older adults*. PLoS One 2014;9:e107541.
- 20 Gopinath B, Sue CM, Kifley A, et al. *The association between olfactory impairment and total mortality in older adults*. J Gerontol A Biol Sci Med Sci 2012;67:204-9.
- 21 Swan GE, Carmelli D. *Impaired olfaction predicts cognitive decline in non-demented older adults*. Neuroepidemiology 2002;2:58-67.
- 22 Sivam A, Wroblewski KE, Alkorta-Aranburu G, et al. *Olfactory dysfunction in older adults is associated with feelings of depression and loneliness*. Chem Senses 2016;41:293-9.
- 23 Negoius S, Hummel T, Symmonk A, et al. *Olfactory volume predicts therapeutic outcome in major depressive disorder*. Brain Imaging Behav 2016;10:367-72.
- 24 Marine N, Boriana A. *Olfactory markers of depression and Alzheimer's disease*. Neurosci Biobehav Rev 2014;45:262-70.
- 25 Naudin M, Mondon K, El-Hage W, et al. *Long-term odor recognition memory in unipolar major depression and Alzheimer's disease*. Psychiatry Res 2014;220:861-6.
- 26 Ansoleaga B, Garcia-Esparcia P, Llorens F, et al. *Dysregulation of brain olfactory and taste receptors in AD, PSP and CJD, and AD-related model*. Neuroscience 2013;248:369-82.
- 27 Cassano T, Romano A, Macheda T, et al. *Olfactory memory is impaired in a triple transgenic model of Alzheimer disease*. Behav Brain Res 2011;224:408-12.
- 28 Desai M, Agadi JB, Karthik N, et al. *Olfactory abnormalities in temporal lobe epilepsy*. J Clin Neurosci 2015;22:1614-8.
- 29 Kohler CG, Moberg PJ, Gur RE, et al. *Olfactory dysfunction in schizophrenia and temporal lobe epilepsy*. Neuropsychiatry Neuropsychol Behav Neurol 2001;14:83-8.
- 30 Ross GW, Petrovitch H, Abbott RD, et al. *Association of olfactory dysfunction with risk for future Parkinson's disease*. Ann Neurol 2008;63:167-73.
- 31 Siderowf A, Jennings D, Eberly S, et al.; PARS Investigators. *Impaired olfaction and other prodromal features in the Parkinson At-Risk Syndrome Study*. Mov Disord 2012;27:406-12.
- 32 Wehling E, Nuess H, Wollschlaeger D, et al. *Olfactory dysfunction in chronic stroke patients*. BMC Neurol 2015;15:199.
- 33 Aliani M, Udenigwe CC, Girgih AT, et al. *Aroma and taste perceptions with Alzheimer's disease and stroke*. Crit Rev Food Sci Nutr 2013;53:760-9.
- 34 Xydakis MS, Mulligan LP, Smith AB, et al. *Olfactory impairment and traumatic brain injury in blast-injured combat troops: a cohort study*. Neurology 2015;84:1559-67.
- 35 Le Floch JP, Le Lièvre G, Labroue M. *Smell dysfunction and related factors in diabetic patients*. Diabetes Care 1993;16:934-7.
- 36 Pattersen DS, Turner P, Smart JV. *Smell threshold in diabetes mellitus*. Nature 1966;209:625.
- 37 Keverne EB. *The vomeronasal organ*. Science 1999;286:716-20.
- 38 Grazzidei PP. *Cell dynamics in the olfactory mucosa*. Tissue Cell 1973;5:113-31.
- 39 Enwere E, Shingo T, Gregg C, et al. *Aging results in reduced epidermal growth factor receptor signalling, diminished olfactory neurogenesis and deficits in fine olfactory discrimination*. J Neurosci 2004;24:8354-65.
- 40 Ressler KJ, Sullivan SL, Buck LB. *A zonal organisation of receptor gene expression in the olfactory epithelium*. Cell 1993;73:597-609.
- 41 Hanchute NK, Kondoh K, Lu Z, et al. *Single-cell transcriptomics reveals receptor transformations during olfactory neurogenesis*. Science 2015;350:1251-5.
- 42 Xia J, Sellers LA, Oxley D, et al. *Urinary pheromones promote ERK/Akt phosphorylation, regeneration and survival of vomeronasal (V2R) neurons*. Eur J Neurosci 2006;24:3333-42.
- 43 Broad KD, Keverne EB. *The post-natal chemosensory environment induces epigenetic changes in vomeronasal receptor gene expression and a bias in olfactory preference*. Behav Genet 2012;42:461-71.
- 44 Loo AT, Youngentob SL, Kent PF, et al. *The ageing olfactory epithelium: neurogenesis, response to damage, and odorant induced activity*. Int J Dev Neurosci 1996;14:881-900.
- 45 Brann JH, Firestein SJ. *A lifetime of neurogenesis in the olfactory system*. Front Neurosci 2014;8:182.
- 46 Nakamura H, Fujiwara M, Kawasaki M, et al. *Age related changes in dividing cells of the olfactory epithelium of the guinea pig*. Eur Arch Otorhinolaryngol 1998;255:289-92.
- 47 Hirai T, Kojima S, Shimada A, et al. *Age-related changes in the olfactory system of dogs*. Neuropathol Appl Neurobiol 1996;22:531-9.
- 48 Weiler E, McCulloch MA, Farbman AI. *Proliferation in the vomeronasal organ of the rat during post-natal development*. Eur J Neurosci 1999;11:700-11.
- 49 Weiler E. *Postnatal development of the rat vomeronasal organ*. Chem Senses 30 (Suppl 10) ii27-ii28.
- 50 Brann JH, Firestein S. *Regeneration of new neurons is preserved in aged vomeronasal epithelia*. J Neurosci 2010;30:15686-94.
- 51 Guillemot F, Lo LC, Johnson JE, et al. *Mammalian achaete-scute homolog 1 is required for the early development of olfactory and autonomic neurons*. Cell 1993;75:463-76.
- 52 Schwob JE, Youngentob SL, Mezza RC. *Reconstitution of the rat olfactory epithelium after methyl bromide induced lesion*. J Comp Neurol 1995;359:15-37.
- 53 Blanco-Hernandez E, Valle-Leija P, Romosa-Signoret V, et al. *Odor memory stability after reinnervation of the olfactory bulb*. PLoS One 2012;7:e46338.
- 54 Iwema CL, Fang H, Kurtz DB, et al. *Odorant receptor expression patterns are restored in lesion recovered rat olfactory epithelium*. J Neurosci 2004;24:356-69.
- 55 Suzukawa K, Kondo K, Kanaya K, et al. *Age related*

- changes of the regenerative mode in the mouse peripheral olfactory system following olfactotoxic drug methimazole-induced damage. *J Comp Neurol* 2011;519:2154-74.
- 56 Milde S, Adalbert R, Elamen MH, et al. Axonal transport declines with age in two distinct phases separated by a period of relative stability. *Neurobiol Ageing* 2015;36:971-81.
 - 57 Adalbert R, Coleman MP. Review: Axon pathology in age-related neurodegenerative disorders. *Neuropathol Appl Neurobiol* 2013;39:90-108.
 - 58 Millicamps S, Julien JP. Axonal transport deficits and neurodegenerative diseases. *Nat Rev Neurosci* 2013;14:161-75.
 - 59 Benignus VA, Prah JD. Olfaction: anatomy, physiology and behaviour. *Environ Health Perspect* 1982;44:15-21.
 - 60 Whitman MC, Greer CA. Adult neurogenesis and olfactory system. *Prog Neurobiol* 2009;89:162-75.
 - 61 Buschhuter D, Smitka M, Puschmann S, et al. Correlation between olfactory bulb volume and olfactory function. *Neuroimage* 2008;42:498-502.
 - 62 Wang J, You H, Liu JF, et al. Association of olfactory bulb volume and olfactory sulcus depth with olfactory function in patients with Parkinson disease. *AJNR Am J Neuroradiol* 2011;32:677-81.
 - 63 Smith CG. Age incident of atrophy of olfactory nerves in man. *J Comparat Neurol* 1942;77:589-94.
 - 64 Kishikawa M, Iseki M, Nishimura M, et al. A histopathological study on senile changes in the human olfactory bulb. *Acta Pathol Jpn* 1990;40:255-60.
 - 65 Kobayashi R, Takahashi-Fujigasaki J, Shiozawa S, et al. α -Synuclein aggregation in the olfactory bulb of middle aged common marmoset. *Neurosci Res* 2016;106: 55-61.
 - 66 Seib DR, Martin-Villalba A. Neurogenesis in the normal ageing hippocampus; a mini review. *Gerontology* 2015;61:327-35.
 - 67 Shook BA, Manz DH, Peters JJ, et al. Spatiotemporal changes to the subventricular zone stem cell pool through aging. *J Neurosci* 2012;32:6947-56.
 - 68 Farbman AI. *Cell biology of olfaction*. Cambridge: Cambridge University Press 1992.
 - 69 Bekkers JM, Suzuki N. Neurons and circuits for odor processing in the piriform cortex. *Trends Neurosci* 2013;36:429-38.
 - 70 Kilpatrick L, Cahill L. Modulation of memory consolidation for olfactory learning by reversible inactivation of the basolateral amygdala. *Behav Neurosci* 2003;117:184-8.
 - 71 Meurisse M, Chaillou E, Lévy F. Afferent and efferent connections of the cortical and medial nuclei of the amygdala in sheep. *J Chem Neuroanat* 2009;37:87-97.
 - 72 Keller M, Perrin G, Meurisse M, et al. Cortical and medial amygdala are both involved in the formation of olfactory offspring memory in sheep. *Eur J Neurosci* 2004;20: 3433-41.
 - 73 De La Rosa-Prieto C, De Moya-Pinilla M, Saiz-Sanchez D, et al. Olfactory and cortical projections to bulbar and hippocampal adult-born neurons. *Front Neuroanat* 2015;9:4.
 - 74 Petrusis A, Alvarez P, Eichenbaum H. Neural correlates of social odor recognition of individual distinctive social odours within entorhinal cortex and ventral subiculum. *Neuroscience* 2005;130:259-74.
 - 75 Alvarez P, Lipton PA, Melrose R, et al. Differential effects of damage within the hippocampal region on memory for a natural, nonspatial Odor-Odor Association. *Learn Mem* 2001;8:79-86.
 - 76 Alvarez P, Eichenbaum H. Representations of odors in the rat orbitofrontal cortex change during and after learning. *Behav Neurosci* 2002;116:421-33.
 - 77 Broad KD, Hinton MR, Keverne EB, et al. Involvement of the medial prefrontal cortex in mediating behavioural responses to odour cues rather than olfactory recognition memory. *Neuroscience* 2002;114:715-29.
 - 78 Segura B, Baggio HC, Solana E, et al. Neuroanatomical correlates of olfactory loss in normal aged subjects. *Behav Brain Res* 2013;246:148-53.
 - 79 Adjei S, Houck AL, Ma K, et al. Age dependent alterations in the number, volume, and localization of islands of Calleja within the olfactory tubercle. *Neurobiol Aging* 2013;34:2676-82.
 - 80 Chapman TR, Barrientos RM, Ahrendsen JT, et al. Aging and infection reduce the expression of specific brain-derived neurotrophic factor mRNAs in hippocampus. *Neurobiol Aging* 2012;33:832.
 - 81 Bedard A, Parent A. Evidence of newly generated neurons in the human olfactory bulb. *Brain Res Dev Brain Res* 2004;151:159-68.
 - 82 De La Rosa-Prieto C, De Moya-Pinilla M, Saiz-Sanchez D, et al. Olfactory and cortical projections to bulbar and hippocampal adult-born neurons. *Front Neuroanat* 2015;9:4.
 - 83 Carrier N, Kabbaj M. Sex differences in the antidepressant-like effects of ketamine. *Neuropharmacology* 2013;70:27-34.
 - 84 Kempermann G. *Adult neurogenesis 2*. New York, NY: Oxford University Press, 2013.
 - 85 Chae CH, Jung SL, An SH, et al. Swimming exercise stimulates neurogenesis in the subventricular zone via increase in synapsin I and nerve growth factor levels. *Biol Sport* 2014;31:309-14.
 - 86 Brown J, Cooper-Kuhn CM, Kempermann G, et al. Enriched environment and physical activity stimulate hippocampal but not olfactory bulb neurogenesis. *Eur J Neurosci* 2003;17:2042-6.
 - 87 Bekinschtein P, Oomen CA, Saksida LM, et al. Effects of environmental enrichment and voluntary exercise on neurogenesis, learning and memory, and pattern separation: BDNF as a critical variable? *Semin Cell Dev Biol* 2011;22:536-42.
 - 88 Rothman S, Mattson M. Activity-dependent, stress-responsive BDNF signaling and the quest for optimal brain health and resilience throughout the lifespan. *Neuroscience* 2013;239:228-40.
 - 89 Llorens-Martin M, Torres-Alemán I, Trejo JL. Growth factors as mediators of exercise actions on the brain. *Neuromol Med* 2008;10:99-107.
 - 90 Trejo JL, Llorens-Martin M, Torres-Alemán I. The effects of exercise on spatial learning and anxiety-like behavior are

- mediated by an IGF-I-dependent mechanism related to hippocampal neurogenesis. *Mol Cell Neurosci* 2008;37:402-11.
- ⁹¹ Fabel K, Fabel K, Tam B, et al. *VEGF is necessary for exercise-induced adult hippocampal neurogenesis*. *Eur J Neurosci* 2003;18:2803-12.
 - ⁹² Bayod S, Menella I, Sanchez-Roige S, et al. *Wnt pathway regulation by long-term moderate exercise in rat hippocampus*. *Brain Res* 2014;1543:38-48.
 - ⁹³ Chen M, Do H. *Wnt signalling in neurogenesis during aging and physical activity*. *Brain Sci* 2012;2:745-68.
 - ⁹⁴ Opendak M, Gould E. *Adult neurogenesis: a substrate for experience-dependent change*. *Trends Cogn Sci* 2015;19:151-61.
 - ⁹⁵ Schubert CR, Cruickshanks KJ, Nondahl DM, et al. *Association of exercise with lower long-term risk of olfactory impairment in older adults*. *JAMA Otolaryngol Head Neck Surg* 2013;139:1061-6.
 - ⁹⁶ Monteiro BM, Moreira FA, Massensini AR, et al. *Enriched environment increases neurogenesis and improves social memory persistence in socially isolated adult mice*. *Hippocampus* 2014;24:239-48.
 - ⁹⁷ Bonzano S, Bovetti S, Fasolo A, et al. *Odour enrichment increases adult-born dopaminergic neurons in the mouse olfactory bulb*. *Eur J Neurosci* 2014;40:3450-7.
 - ⁹⁸ Rochefort C, Gheusi G, Vincent JD, et al. *Enriched odor exposure increases the number of newborn neurons in the adult olfactory bulb and improves odor memory*. *J Neurosci* 2002;22:2679-89.
 - ⁹⁹ Mandaïron N, Stack C, Kiselycznyk C, et al. *Enrichment to odors improves olfactory discrimination in adult rats*. *Behav Neurosci* 2006;120:173-9.
 - ¹⁰⁰ Escanilla O, Mandaïron N, Linster C. *Odor-reward learning and enrichment have similar effects on odor perception*. *Physiol Behav* 2008;94:621-6.
 - ¹⁰¹ Richardson BE, Vander Woude EA, Sudan R, et al. *Altered olfactory acuity in the morbidly obese*. *Obesity Surgery* 2004;14:967-9.
 - ¹⁰² Patel ZM, Del Gaudio JM, Wise SK. *Higher body mass index is associated with subjective olfactory dysfunction*. *Behav Neurol* 2015;2015:675635.
 - ¹⁰³ Warner-Schmidt JL, Duman RS. *Hippocampal neurogenesis: opposing effects of stress and antidepressant treatment*. *Hippocampus* 2006;16:239-49.
 - ¹⁰⁴ Siopi E, Denizet M, Gabellec MM, et al. *Anxiety- and depression-like states lead to pronounced olfactory deficits and impaired adult neurogenesis in mice*. *J Neurosci* 2016;36:518-31.
 - ¹⁰⁵ Croy I, Symmank A, Schellong J, et al. *Olfaction as a marker for depression in humans*. *J Affect Disord* 2014;160:80-6.
 - ¹⁰⁶ Greiner RS, Moriguchi T, Slotnick BM, et al. *Olfactory discrimination deficits in n-3 fatty acid-deficient rats*. *Physiol Behav* 2001;72:379-85.
 - ¹⁰⁷ Beltz BS, Tlusty MF, Benton JL, et al. *Omega-3 fatty acids upregulate adult neurogenesis*. *Neurosci Lett* 2007;415:154-8.
 - ¹⁰⁸ Gopinath B, Sue CM, Flood VM, et al. *Dietary intakes of fats, fish and nuts and olfactory impairment in older adults*. *Br J Nutr* 2015;114:240-7.
 - ¹⁰⁹ Dong S, Zeng Q, Mitchell ES, et al. *Curcumin enhances neurogenesis and cognition in aged rats: implications for transcriptional interactions related to growth and synaptic plasticity*. *PLoS One* 2012;7:e31211.
 - ¹¹⁰ An L, Zhang YZ, Yu NJ, et al. *The total flavonoids extracted from Xiaobuxin-Tang up-regulate the decreased hippocampal neurogenesis and neurotrophic molecules expression in chronically stressed rats*. *Prog Neuropsychopharmacol Biol Psychiatry* 2008;32:1484-90.
 - ¹¹¹ Casadesus G, Shukitt-Hale B, Stellwagen HM, et al. *Modulation of hippocampal plasticity and cognitive behavior by short-term blueberry supplementation in aged rats*. *Nutr Neurosci* 2004;7:309-16.
 - ¹¹² Bonnet E, Touyarot K, Alfors S, et al. *Retinoic acid restores adult hippocampal neurogenesis and reverses spatial memory deficit in vitamin A-deprived rats*. *PLoS One* 2008;3:e3487.
 - ¹¹³ Han ME, Park KH, Baek SY, et al. *Inhibitory effects of caffeine on hippocampal neurogenesis and function*. *Biochem Biophys Res Commun* 2007;356:976-80.
 - ¹¹⁴ Ogawa T, Takezawa K, Shimizu S, et al. *Valproic acid promotes neural regeneration of olfactory epithelium in adult mice after methimazole-induced damage*. *Am J Rhinol Allergy* 2014;28:e95-9.
 - ¹¹⁵ Xia J, Broad KD, Emson PC, et al. *Epigenetic modification of vomeronasal (V2r) precursor neurons by histone deacetylation*. *Neuroscience* 2010;169:1462-72.
 - ¹¹⁶ Inta D, Lima-Ojeda JM, Lau T, et al. *Electroconvulsive therapy induces neurogenesis in frontal rat brain areas*. *PLoS One* 2013;8:e69869.
 - ¹¹⁷ Henkin RI, Potolicchio SJ Jr, Levy LM. *Improvement in smell and taste dysfunction after repetitive transcranial magnetic stimulation*. *Am J Otolaryngol* 2011;32:38-46.
 - ¹¹⁸ Ueyama E, Ukai S, Ogawa A, et al. *Chronic repetitive transcranial magnetic stimulation increases hippocampal neurogenesis in rats*. *Psychiatry Clin Neurosci* 2011;65:77-81.
 - ¹¹⁹ Arias-Carrión O, Verdugo-Díaz L, Fera-Velasco A, et al. *Neurogenesis in the subventricular zone following transcranial magnetic field stimulation and nigrostriatal lesions*. *J Neurosci Res* 2004;78:16-28.